

 	HOSPITAL SULTAN ABDUL AZIZ SHAH UNIVERSITI PUTRA MALAYSIA Document Code: HSAAS/OFTAL/BR160	<i>Patient Sticker</i>
	OT / PROCEDURE FORM	

DATE		PROCEDURE	
VITAL SIGNS	BP :	HR :	DXT :

Procedure steps:

1. Patient in supine position comfortably. Alert and conscious. Pink.
2. Dilate the pupil. Topical anaesthesia was applied to the affected RE/LE.
3. 10% povidone iodine solution was applied to the periocular skin, lids, eyelashes.
4. Sterile drape was placed to the RE/LE.
5. Sterile lid speculum was placed to the RE/LE.
6. 5% povidone ophthalmic iodine solution was applied to the RE using 3cc syringe to the conjunctival sac, and wait for 90 seconds.
7. Rinse the eye with ophthalmic saline solution.
8. Direct the patient to look away from the injection site. Mark an injection site at an area 4.0mm/3.5mm Superotemporally/.....) posterior to the limbus, avoiding the horizontal meridian and aiming toward the center of the globe.
9. Slowly deliver the injection volume, then the needle was withdrawn gently.
10. Gutt CMC/.....) was applied topically to the eye.
11. Globe was felt, not hard. Gross visual acuity was tested using fingers.

Intraoperative complications:

- 1.
- 2.

Plan:

1. Continue Gutt Chloramphenicol QID for 1/52
2. TCA 1/12 for Visual acuity, DBE or TCA STAT (to nearby hospital) if develop symptoms of redness, swollen eyes, reduced vision, eye discharge, eye pain, floaters, flashes

Performed by,

.....

Supervised by,

.....